

This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.




Prentox®

ExciteR™

With Pyrethrins,
a Floral Insecticide

Contains 6% Py and 60% PBO.

- *Powered by Pyronyl*
- *Rapid knock down and kill*
- *Powerful flushing action*
- *A tank mix partner for resistance management*
- *Basic component for flea programs*



ACTIVE INGREDIENTS:

Pyrethrins	6.0%
Piperonyl Butoxide, Technical*	60.0%
OTHER INGREDIENTS**:	34.0%
TOTAL:	100.0%

*Equivalent to min. 48.0% (butylcarbityl)(6-propypiperonyl) ether and 12.0% related compounds.
 **Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside label booklet for First Aid, Precautionary Statements
 and Directions for Use including Storage and Disposal instructions
 For Chemical Spill, Leak, Fire, or Exposure, Call: CHEMTREC 1-800-424-9300
 For Medical Emergencies Only, Call: 866-257-4118



EPA Reg. No. 655-798

Manufactured by:
 Prentiss LLC
 501 Cascade Pointe Lane, Suite 103
 Cary, NC 27513

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
If swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center, physician, or PROSAR at 866-257-4118 for treatment advice. • Do not induce vomiting unless told to do so by the Poison Control Center or physician. • Do not give any liquid to the person. • Avoid alcohol. • Do not give anything by mouth to an unconscious or convulsing person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center, physician, or PROSAR at 866-257-4118 for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center, physician, or PROSAR at 866-257-4118 for treatment advice.
If inhaled;	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth. • Call a Poison Control Center, physician, or PROSAR at 866-257-4118 for treatment advice.
NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 866-257-4118.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or inhaled. Causes eyeirritation. Avoid breathing spray mist. Avoid contact with eyes. Wear protective clothing. Wash skin thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. Do not apply as a space spray while food processing is underway. Food processing surfaces and equipment must be covered during treatment or cleaned with an effective detergent and rinsed with potable water before reuse. Except in Federally inspected meat and poultry plants, food processing operations may continue when this product is applied as a surface spray with care and in accordance with the directions and precautions given above.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crab and other aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high watermark. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Tank Mixes in Combination With Residual Insecticides: This product can be mixed with Cynoff®² EC Insecticide #279-3081, Cynoff®² WP Insecticide #279-3070, Conquer®³ Residual Insecticide Concentrate # 1021-1565-57076, Prelude®⁴ Termiticide/Insecticide #10182-95, Demon®⁴ EC Insecticide #100-1004, Perm-Up®⁵ 3.2 EC #70506-9, Astro®² Insecticide #279-3141, LG⁶ Permethrin 3.2 Termiticide/Insecticide #71532-2, Phaeton Corporation 354, Atroban®⁸ 11% EC #773-59, Prentox®⁹ Prenbay™⁹ 1.5 EC #655-796, Prentox®⁹ Perm-X™⁹ 1E #655-783, at the rate of ¼ to ½ fl. ounce (equivalent to ½ to 1 tablespoon or 7.4 to 14.8 ml) per gallon of finished spray, for use as a flushing and knockdown agent to aid in

control of crawling insects and flying insects as listed on the labels of each product, in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. To provide increased control of resistant German Cockroaches: tank mix 2.125 to 4.25 fl. ounces of ExciteR™ with the appropriate amount of companion insecticide label. Prior to tank mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. When tank-mixing ExciteR with wettable powder formulations, first add the wettable powder to the diluent and agitate to thoroughly suspend it in solution and then add the desired amount of ExciteR to the diluted wettable powder. Tank –mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be tank-mixed with any product with label prohibitions against such mixing.

Trademark designations: ¹McLaughlin Gormley King Co., ²FMC Company, ⁴Syngenta Professional Products, ⁵United Phosphorus, ⁶LG Chem Investment, Inc., ⁷Phaeton Corporation, ⁸Intervet Inc., ⁹Prentiss Incorporated.

A highly versatile, synergized, pyrethrin, insecticide concentrate for indoor or outdoor application as a surface, space, area or contact spray. ExciteR may be applied through mechanical, aerosol or ULV generators as well as thermal fogging equipment and conventional fogging or spraying equipment. ExciteR may be applied as a general surface, space, spot or crack and crevice treatment in both food and non-food areas of food handling establishments and is authorized for use in all areas of official establishments operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs.

ExciteR is intended for use in and around buildings and structures as well as modes of transport. **INDOOR USE AREAS:** Apartment buildings, bakeries, beverage plants, bottling facilities, breweries, cafeterias, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, flour mills, food processing plants, frozen food plants, grain elevators, grain mills, granaries, homes, hospitals, hotels, houses, industrial buildings, kennels, kitchens, laboratories, manufacturing facilities, mausoleums, meat processing and packaging plants, meat and vegetable canneries, motels, nursing homes, office buildings, restaurants, schools, spice plants, stores, supermarkets, warehouses, wineries and similar structures. **OUTDOOR AREAS:** Drive-in restaurants, drive-in theaters, golf courses, parks, playgrounds, recreational areas, urban areas and zoos. **MODES OF TRANSPORT:** Aircraft (do not use in occupied aircraft cabins), buses, trucks, trailers, rail cars and marine vessels.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

ExciteR may be diluted with water or oil and applied with conventional application equipment including, but not limited to, compressed air sprayers, power sprayers, mechanic alaerosol generators (ULV generators, or cold foggers) and thermal foggers (oil dilutions only). When diluting with water, prepare only the amount needed for immediate use and agitate occasionally during use. Holding a water dilution more than 24 hours is not recommended. For oil dilutions that are applied in food handling establishments, a deodorized petroleum oil which conforms with CFR 21 (especially section 172.884) is required (such as Exxon Isopar M, Ashland LowOdor Base Solvent or the equivalent).

Fluid Ounces of ExciteR Per Gallon	% Pyrethrins	Approximate % Piperonyl Butoxide
0.25	0.0125	0.125
0.5	0.025	0.25
1.0	0.05	0.5
2.125	0.1	1.0
4.25	0.2	2.0

IN USDA INSPECTED FACILITIES: To control accessible, exposed stages of crawling insects including, but not limited to, *Ants, Cockroaches, Cadelles, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Spider Beetles, Yellow Mealworms*, dilute 2.125 fluid ounces of ExciteR per gallon of water and apply at the rate of 1 gallon to 750 square feet. Thoroughly spray the surfaces where these insects are usually found including walls, baseboards, moldings, doors, cabinets, shelves, sinks and pipes with special attention to cracks and crevices, niches, dark corners, drains and other harborage sites.

FLEA AND TICK CONTROL IN BUILDINGS: Dilute 2.125 to 4.25 fluid ounces of ExciteR per gallon of water and apply the spray thoroughly to infested areas such as pet beds and resting quarters, nearby cracks and crevices, rugs and floor coverings (including under their edges) between and under the cushions of upholstered furniture and other areas where these pests may be present. Entire rugs and furniture should be treated at the rate of 1 gallon of dilute spray per 750 square feet of surface. In combination with conventional residual insecticides: To provide quick knockdown of adult fleas, tank-mix 0.5 to 1.0 fluid ounce of ExciteR with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide label.

In combination with insect Growth Regulators: To provide quick knockdown of adult fleas, tank-mix 2.125 to 4.25 fluid ounces of ExciteR with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide label.

Pet bedding should be removed and replaced with fresh bedding after treatment of pet areas. The pets should also be treated with an EPA registered product before they are allowed to enter the treated area.

PANTRY PEST CONTROL: To control exposed adult and immature stages of Stored Product Insects including, but not limited to, *Almond Moths, Angoumois Grain Moths, Cadelles, Cigarette Beetles, Confused Flour Beetles, Drugstore Beetles, Flat Grain Beetles, Granary Weevils, Indianmeal Moths, Lesser Grain Borers, Maize Weevils, Mediterranean Flour Moths, Merchant Grain Beetles, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles and Squarenecked Grain Beetles*, dilute 2.125 to 4.25 fluid ounces of ExciteR per gallon of water and apply to cupboards, shelves and storage areas. Remove all utensils, unprotected foodstuffs (or any with original packaging that has been opened), shelf paper and other objects before treatment. Allow treated surfaces to dry and then cover shelves with clean paper before replacing any utensils, foodstuffs or other items. Discard any foodstuff that was accidentally contaminated with the spray solution.

PREMISE PEST CONTROL: Termite swarmers: To control swarming termites, dilute 2.125 to 4.25 fluid ounces of ExciteR per gallon of water and apply as a surface spray at the rate of 1 gallon per 750 square feet. Apply as a crack and crevice treatment to the locations from which the termites are emerging.

Carpet Beetles: To control carpet beetles (black, furniture and varied), dilute 4.25 fluid ounces of ExciteR per gallon of water and apply to the entire carpet at the rate of 1 gallon per 750 square feet. Make localized applications under the edge of carpets, floor coverings, and furniture as well as along baseboards, in closets and to infested areas of shelving.

BED BUGS: To control *Bed bugs*, dilute 2.125 fluid ounces of ExciteR per gallon of water and apply thoroughly to the top, ends, sides and bottoms of mattresses paying special attention to tufts, seams and folds. Treat bedsprings and frames, including all cracks and joints. Apply as a crack and crevice treatment to all baseboards, moldings, flooring, closets, shelves, furniture and picture frames that may provide harborage to bedbugs.

WEBBING CLOTHES MOTHS: To control *Clothes Moths*, remove infested articles from storage, brush them thoroughly and air-dry them in sunlight for several hours. Dilute 2.125 fluid ounces of ExciteR per gallon of water and apply as a crack and crevice treatment to empty chests, bureaus and storage sites. After airing, infested articles may be lightly treated with the spray solution (keep the nozzle at least three feet from the fabric to avoid staining).

FOR INDOOR APPLICATIONS: Remove pets (including birds) from the area being treated and cover fish aquaria before spraying.

INDOOR USE

To control crawling and flying insects in sites that include, but are not limited to, food processing plants, (INCLUDING USDA INSPECTED FACILITIES), industrial installations and warehouses, ExciteR may be diluted with water and applied as a space spray. For best results, close doors and windows before spraying and keep them closed for 30 minutes after treatment. The applicator must vacate the area after treatment and ventilate before reoccupying. Where oil residues are not undesirable, ExciteR can be diluted in deodorized base oil instead of water and applied with mechanical, thermal or ULV applicators.

The use of this product, as a space spray, in food processing or food handling establishments must be confined to time periods when the plant is not in operation. Foods must be covered or removed during treatment. Food processing surfaces and equipment must be covered during treatment or cleaned with an effective detergent and rinsed with potable water before use.

CRAWLING AND FLYING INSECTS: For control of accessible, exposed stages of CRAWLING INSECTS including, but not limited to, *Ants, Cockroaches, Cadelles, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Spider Beetles, Yellow Mealworms and Flying Insects* including, but not limited to, *Angoumois Grain Moths, Cheese Skippers, Fruit Flies, Fungus Gnats, Gnats, House Flies, Indianmeal Moths, Mosquitoes, Mediterranean Flour Moths, Small Flying Moths, Tobacco Moths*, dilute 10.67 fluid ounces of ExciteR per gallon of water or oil and apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray towards the ceiling and upper corners of the area and behind obstructions. Vacate the treated area and keep the area closed for at least 30 minutes after treatment. Ventilate the area before reoccupying. Repeat treatment as necessary.

FLYING INSECTS: For control of accessible, exposed stages of flying insects including, but not limited to, *Angoumois Grain Moths, Cheese Skippers, Fruit Flies, Fungus Gnats, Gnats, House Flies, Indianmeal Moths, Mosquitoes, Mediterranean Flour Moths, Small Flying Moths, Tobacco Moths*, dilute 2.67 fluid ounces of ExciteR per gallon of water or oil and apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray towards the ceiling and upper corners of the area and behind obstructions. Vacate the treated area and keep the area closed for at least 30 minutes after treatment. Ventilate the area before reoccupying. Repeat treatment as necessary.

VOID AREA TREATMENT: Undiluted ULV treatment: To treat inaccessible voids including, but not limited to, attics, false ceilings, wall and equipment voids, calculate the volume of the void and apply ExciteR through ULV equipment adjusted for crack and crevice application at the rate of 0.33 fluid ounces per 1000 cubic feet of space.

OUTDOOR USE

PERIMETER TREATMENTS: To control *Ants, Bees, Centipedes, Cockroaches, Crickets, Firebrats, Fleas, Ground Beetles, Millipedes, Pillbugs, Silverfish, Sowbugs, Spiders, Ticks and Wasps*, dilute 2.125 to 4.25 fluid ounces of ExciteR per gallon of water and apply at the rate of 1 gallon per 750 square feet to a band of soil and vegetation 6 to 10 feet wide around and adjacent to buildings. Treat the building foundation to a height of 2 to 3 feet at the rate of 1 gallon per 750 square feet.

TICK CONTROL: To control *Brown Dog Ticks* around residential areas, rural fields, woods, lots, roadsides, pathways and coastal vegetation areas, dilute 4.25 fluid ounces of ExciteR per gallon of water and apply with a mist blower (or other similar equipment) calibrated to deliver 3 gallons per acre. Apply during the peak of infestation and repeat as necessary to maintain control. Thorough applications should be made to the vegetation upon which the ticks rest.

FLEA CONTROL: To control fleas around residential areas, dog houses, kennels, paths, patios and under plants, shrubs, trees or other shaded, moist areas where pets may rest, dilute 4.25 fluid ounces of ExciteR per gallon of water and apply at the rate of 1 gallon per 750 square feet.

In combination with other insecticides: To provide quick knockdown of insects when used with a residual insecticide, tank-mix 1.0 to 2.125 fluid ounces of ExciteR with the proper amount of companion insecticide and apply per the Directions for Use on the companion insecticide label.

USE ON TURF AND GRASS: Used alone as a contact spray: To control *Ants, Armyworms, Billbugs, Chinch Bugs, Chiggers, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Hyperodes Weevils (adults), Japanese Beetles*

(adults), Mole Crickets, Sod Webworms And Ticks, dilute and apply per the instructions in the following table:

Treatment Area (Square Feet)	Fluid Ounces of ExciteR	Suggested Volume^a Of Water (Gallons)
1000	0.25 to 0.5	2.5 to 5.0
5000	1.25 to 2.5	12.5 to 25.0
20000	5.0 to 10.0	50.0 to 100.0
43560 ^b	12.0 to 24.0	110.0 to 220.0

^a Dilute with enough water to obtain thorough coverage.

^b 43560 square feet = 1 acre.

In combination with other insecticides: To provide quick knockdown of insects when used with a residual insecticide, tank-mix ExciteR at the rate listed above with the proper amount of companion insecticide and apply per the Directions for Use on the companion insecticide label.

As a turf pest diagnostic aid: To detect turf insects prior to making an insecticide application or to evaluate control from previous treatments, dilute 0.5 fluid ounces of ExciteR per gallon of water and apply evenly with a sprinkling can over one square yard of turf. Record the species and number of insects present ten minutes after application. Sample 3 to 5 sites per 5000 square feet. Note: This procedure does not bring white grubs or billbug grubs to the surface. Use other methods to sample for these pests.

PETS

PRE-TREATMENT DIRECTIONS

READ ENTIRE LABEL BEFORE EACH USE USE ONLY ON DOGS TO KILL FLEAS AND TICKS: Do not use on dogs under 12 weeks.

SPECIAL ANIMALS

Consult a veterinarian or call the National Pesticide Information Center at 1-800-858-7378, before using this product on debilitated, aged, pregnant nursing or medicated animals.

ADVERSE REACTION INFORMATION

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your dog with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately, or call the National Pesticide Information Center at 1-800-858-7378.

APPLICATION DIRECTIONS

TO CONTROL FLEAS AND TICKS ON DOGS: Dogs may be treated with ExciteR by diluting at the rate of 1.0 to 2.125 fluid ounces per gallon of water and spraying or sponging thoroughly (to penetrate the hair and contact the skin) to all parts of the animal's body except the eyes, nose, mouth and genitalia. For best results against fleas and ticks on dogs, the animal's quarters should also be treated and bedding should be removed and replaced after the pet areas are treated.

REAPPLICATION DIRECTIONS

Do not reapply product for 7 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Clean container promptly after emptying. {Containers 5 gallons or less} Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate. {Containers larger than 5 gallons} Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Direction for Use of this Product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prentiss LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prentiss LLC and Seller harmless for any claims relating to such factors.

Prentiss LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prentiss LLC, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, PRENTISS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law Prentiss LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRENTISS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRENTISS LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prentiss LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prentiss LLC.

PRENTOX® - Registered Trademark of Prentiss LLC

ExciteR™ - Trademark of Prentiss LLC